

WO9511974

Publication Title:

HUMAN PEROXISOME PROLIFERATOR ACTIVATED RECEPTOR

Abstract:

Abstract of WO9511974

A human peroxisome proliferation activated receptor gene is purified from the environment in which it naturally occurs, and preferably provided within an expression vector.

Data supplied from the esp@cenet database - Worldwide

Courtesy of <http://v3.espacenet.com>



INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification ⁶ : C12N 15/12, C07K 14/705		A2	(11) International Publication Number: WO 95/11974
			(43) International Publication Date: 4 May 1995 (04.05.95)
(21) International Application Number: PCT/US94/11897 (22) International Filing Date: 19 October 1994 (19.10.94) (30) Priority Data: 08/141,500 22 October 1993 (22.10.93) US 08/143,215 26 October 1993 (26.10.93) US (71) Applicant: LIGAND PHARMACEUTICALS, INC. [US/US]; 9393 Towne Center Drive, San Diego, CA 92121 (US). (72) Inventor: MUKHERJEE, Ranjan; 11341 Avenida De Los Lobos, San Diego, CA 92127 (US). (74) Agents: CHEN, Anthony, C. et al.; Lyon & Lyon, First Interstate World Center, Suite 4700, 633 West Fifth Street, Los Angeles, CA 90071-2066 (US).		(81) Designated States: AM, AT, AU, BB, BG, BR, BY, CA, CH, CN, CZ, DE, DK, EE, ES, FI, GB, GE, HU, JP, KE, KG, KP, KR, KZ, LK, LR, LT, LU, LV, MD, MG, MN, MW, NL, NO, NZ, PL, PT, RO, RU, SD, SE, SI, SK, TJ, TT, UA, UZ, VN, European patent (AT, BE, CH, DE, DK, ES, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, ML, MR, NE, SN, TD, TG), ARIPO patent (KE, MW, SD, SZ). Published Without international search report and to be republished upon receipt of that report.	

(54) Title: HUMAN PEROXISOME PROLIFERATOR ACTIVATED RECEPTOR

```

10      20      30      40      50      60      70      80      90      100
1234567890 1234567890 1234567890 1234567890 1234567890 1234567890 1234567890 1234567890 1234567890 1234567890
ATGCTGACAA GGAAGAGCC ACTCTGCCC CTCTGCCC TCGAGCGCG GATCTAGAG AGCCGTTAT CTGAGAGAT CTGCAAGAA ATGGGAACA
HVD TESP LCP LSPLEAG DLE SPLS EEF LQE HGH I
TCCAGAGAT TTGCGATCC ATGGGCGAG ATAGTTCTGG AAGCTTTGGC TTACGGAAT ACCAGTATT AGGAGCTGT CTGCGCTAG ATGCGCTGCT 200
QEI SQS IGE DSS G SFG FTEY QYL GSC PGS D GSV
CATCAGGAC AGCTTTTCAE CAGCTTOSAG CCGCTGCTCG GTGACTTATE CTGTGCTCC CGCAGCGTG GAGAGCTTC CAGTGGAGC ATTGAACATC 300
T T O T L S P A S S P S S V T Y P V V P G S V D E S P S G A L N I
GAATGTAGAA TGTGCGGGA CAGGCGTCA GGTATCATY ACGGAGTCA GCGGTGTAA GCGTCAAGG GCTTCTTGG GCGAAGGAT GCACTCAAGC 400
E C R I C G D K A S G Y H Y G V H A C E G C K G F F R R T I R L K L
TGGTGTATGA CAGTGGGAC CGCAGCTGCA AGATCCAGAA AAGAAGAGA AACAAATGCC AGTATTGTGG ATTTCACAG TGGCTTTCTG TCGGATGTC 500
V Y D K C D R S C K I Q K K N R N K C Q Y C R F H K C L S V G M S
ACACAAGCG ATGTTGTTT GAGGAATGCC AAGATCTGAG AAGCAAAAC TGAAGCAGA AATCTTACC TGTGAACATG ACATAGAAGA TTCTGAAC 600
H N A I R F G R H P R S E K A K L K A E I L T C E H D I E D S E T
GCAGATCTCA AATCTCTGC CAGAGAATC TACGAGGCT ACTTGAGAA CTTCACATG AACAGGTCA AAGCGCGGT CATCTCTCA GGAAGGCCA 700
A D L K S L A K R I Y E A Y L K N F H M N K V K A R Y I L S G K A S
GTAACAATCC AGCTTTTCT ATACATGTA TGGAGACACT GTGTATGCT GAGAAGGCC TGTGCGCAA GCTGTGGCC AATGGCATCC AGAACAAGA 800
N H P P F V I H D H E T L C H A E X T L V A K L V A N G I Q N K E
GGCGAGGTC GCACTCTTC ACTGCTGCA GTGAGGTCA GTGAGAGCG TCAAGGAGT CAGGGAATC GCGAAGGCC TCCAGGCTT CCAACAATCT 900
A E V R I F H C C Q C T S V E T V Y E L T E F A K A I P G F A N L
GAGCTGAAGC ATCAAGTAC ATTGCTAATA TACGAGTIT ATGAGGCAAT ATGCGCATG CTGTCTCTG TGAAGACAA AGACGGATG CTGTAGGCT 1000
D L N D Q V T L L K Y G V Y E A I F A H L S S V H N K D G M L V A Y
ATGGAATGG GTTTAATCT GTGGAATCC TAAAGGCT AGGAAGCG TTCTGTSATA TCATGAGGC CAGTTTAT TTTGCGTGA AGTTCAATGC 1100
G N G F I T R E F L K S L R K P F C D I H E P K F D F A H K F N A
ACTGGAATG GATGACAGT ATATCTCCT TTTTGTGCT GCTATCATY GTGTGAGAA TGTGCTGCG CTCTTAAGC TAGGACAT TGAAGAAATG 1200
L E L D D S D I S L F V A A I T C G D R P G L L N V G H I E K H
CAGGAGGTA TTGTACATGT GCTCAGACT CAGCTGAGA GCAAGCACC GCAGATATC TTCTCTTCC CAAGCTTCT TGAAGAAATG GCAAGCTCC 1300
Q E G I V H V L R L H L Q S H H P D I F L F P K L L Q K H A D L R
GCAAGTGT GAGGAGCAT GGCAGCTGG TGCAGATCAT CAGAAGAGC GAGTGGATG CTGCGCTGCA CCGCTACTG CAGGAGATCT ACAGGAGCAT 1400
Q L V T E H A Q L V Q I I K K T E S D A A L H P L L Q E I Y R D H
GTACTGA
Y X

```

(57) Abstract

A human peroxisome proliferation activated receptor gene is purified from the environment in which it naturally occurs, and preferably provided within an expression vector.

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AT	Austria	GB	United Kingdom	MR	Mauritania
AU	Australia	GE	Georgia	MW	Malawi
BB	Barbados	GN	Guinea	NE	Niger
BE	Belgium	GR	Greece	NL	Netherlands
BF	Burkina Faso	HU	Hungary	NO	Norway
BG	Bulgaria	IE	Ireland	NZ	New Zealand
BJ	Benin	IT	Italy	PL	Poland
BR	Brazil	JP	Japan	PT	Portugal
BY	Belarus	KE	Kenya	RO	Romania
CA	Canada	KG	Kyrgyzstan	RU	Russian Federation
CF	Central African Republic	KP	Democratic People's Republic of Korea	SD	Sudan
CG	Congo	KR	Republic of Korea	SE	Sweden
CH	Switzerland	KZ	Kazakhstan	SI	Slovenia
CI	Côte d'Ivoire	LI	Liechtenstein	SK	Slovakia
CM	Cameroon	LK	Sri Lanka	SN	Senegal
CN	China	LV	Latvia	TD	Chad
CS	Czechoslovakia	LU	Luxembourg	TG	Togo
CZ	Czech Republic	MG	Madagascar	TJ	Tajikistan
DE	Germany	ML	Mali	TT	Trinidad and Tobago
DK	Denmark	MN	Mongolia	UA	Ukraine
ES	Spain			US	United States of America
FI	Finland			UZ	Uzbekistan
FR	France			VN	Viet Nam
GA	Gabon				

DESCRIPTIONHuman Peroxisome Proliferator Activated ReceptorCross Reference to Related Application

This application is a continuation-in-part of Application Docket No. 202/041, titled "Human Peroxisome Proliferator Activated Receptor," filed October 22, 1993, by Mukherjee, the disclosure of which is incorporated herein by reference.

Field of the Invention

This invention relates to the cloning and uses of a human peroxisome proliferator activated receptor.

Background of the Invention

5 A peroxisome proliferator is an agent that induces peroxisomal proliferation. Peroxisome proliferators are a diverse group of chemicals which include unsaturated fatty acids, hypolipidemic drugs, herbicides, leukotriene antagonists, and plasticizers (for a review, see Green,
10 S., 43 Biochem. Pharmacol. 393-400, 1992). Hypolipidemic drugs such as clofibrates have been found to lower triglycerides and cholesterol levels in plasma and to be beneficial in the prevention of ischaemic heart disease in
15 individuals with elevated levels of cholesterol (Havel, R.J. and Kane, J.P., 13 Ann. Rev. Pharmac. 287-308, 1973). Therapeutic use of such drugs, however, is questioned because clofibrates are carcinogens in rats.

Peroxisome proliferator activated receptor (PPAR) is a member of the steroid receptor family. It is activated
20 by peroxisome proliferators. Issemann and Green, 347 Nature 645, 1990, cloned a mouse peroxisome proliferator activated receptor (mPPAR) gene from a mouse liver complementary DNA (cDNA) library. Göttlicher et al., 89 Proc. Nat. Acad. Sci. USA 4653-4657, 1992, cloned a rat
25 peroxisome proliferator activated receptor (rPPAR) gene from a rat liver cDNA library. PPARs from mouse and rat share 97% homology in amino acid sequence and a

particularly well-conserved putative ligand-binding domain. Three members of the *Xenopus* nuclear hormone receptor superfamily have also been found to be structurally and functionally related to the mPPAR (Dreyer et al., 68 Cell 879-887, 1992).

Schmidt et al., 6 Molecular Endocrinology 1634-1641, 1992, cloned a steroid hormone receptor gene, NUC1, from a human osteosarcoma cell cDNA library. The homology between amino acid sequence of NUC1 and that of the mouse PPAR is only 62%. Thus, although it is clear that NUC1 is a member of the PPAR receptor group, it remains to be determined whether NUC1 is the human homolog of the mouse PPAR or a new member of the PPAR family.

Sher et al., 32 Biochemistry 5598-5604, 1993, cloned a human PPAR gene from a human liver cDNA library. This clone has 85% nucleotide sequence homology and 91% amino acid sequence homology with the mPPAR clone.

Summary of the Invention

The present invention relates to the cloning of a human PPAR gene, hPPAR1. The protein encoded by hPPAR1 has 92% homology with the mouse PPAR. It is different from the human PPAR cloned by Sher et al., supra, at two locations in the amino acid sequence, i.e., amino acids 268 and 296.

The hPPAR1 clone can be used for the expression of large amounts of hPPAR1. This human PPAR clone is also useful for screening compounds for improved pharmacological profiles for the treatment of hyperlipidemia with higher potency, efficacy, and fewer side effects. Specifically, the human PPAR clone can be used to screen for compounds active as primary endogenous inducers of the human PPAR. In addition, it is useful for establishing the tissue specific expression pattern of human PPAR. For example, a Northern blot can be used to reveal tissue specific expression of the gene to aid treatment of diseases such as atherosclerosis.

Thus, in a first aspect, the invention features a purified nucleic acid encoding a human PPAR with the nucleotide base sequence shown in Figure 1, and given as SEQ ID NO. 1. By purified nucleic acid is meant that the
5 nucleic acid is separated from its natural environment and from other nucleic acids.

In a second aspect, the present invention features a vector containing the human PPAR gene. This vector may be used for multiplication of the human PPAR gene or
10 expression of the human PPAR gene.

In a preferred embodiment, the vector is an expression vector. In one example, the expression vector is used to make a recombinant human PPAR nucleic acid, which can be used as a specific probe for DNA or RNA
15 complementary to the human PPAR sequence. In another example, the expression vector is used to express human recombinant PPAR protein.

By vector is meant a plasmid or viral DNA molecule into which either a cDNA or a genomic DNA sequence is
20 inserted.

By expression vector is meant a vector that directs protein synthesis from a promoter. In a preferred embodiment, either vector pBacPAK8 (Clontech) or vector pBacPAK9 (Clontech) is used to express the human PPAR in
25 insect cells. In another preferred embodiment, vector pYES2 (Invitrogen) is used to express the human PPAR in yeast cells. In yet another preferred embodiment, pBKCMV (Stratagene) is used to express the human PPAR in mammalian cells.

30 By recombinant human PPAR is meant a non-naturally expressed human PPAR.

Other features and advantages of the invention will be apparent from the following description of the preferred embodiments thereof, and from the claims.

Description of the Preferred Embodiments

Drawings

Figure 1 is the nucleotide and amino acid sequence of hPPAR1; and

5 Figure 2 is a comparison of the amino acid sequences of hPPAR1 and the mouse PPAR.

What follows is an example of the cloning of a human PPAR. Those of ordinary skill in the art will recognize that equivalent procedures can be readily used to isolate
10 human PPAR from cDNA libraries or genomic libraries of other tissues than that exemplified below, namely the liver.

In general, the cloning of the human PPAR involved probing a human liver cell cDNA library with a labeled
15 EcoRI-BglIII fragment (nucleotides 450-909) of the rat PPAR (459 bases). The sequence of the probe is shown in Götlicher et al. supra.

The recipes for buffers, mediums, and solutions in the following examples are given in J. Sambrook, E. F.
20 Fritsch, and T. Maniatis, Molecular Cloning: A Laboratory Manual, 2 Ed., Cold Spring Harbor Laboratory Press, Cold Spring Harbor, New York, 1989.

Example 1: Cloning of a human PPAR

A human PPAR subtype, hPPAR1, was cloned from a human
25 liver 5'-stretch cDNA library (Clontech #HL1115a) in lambda gt10 phages. C600-Hfl coli (Clontech) was grown overnight in LB broth supplemented with 0.2% maltose. A required amount of phage (corresponding to 2 million plaques) was mixed with 200 microliters of 10 mM MgCl₂/10
30 mM CaCl₂ and 1.5 milliliters of the overnight C600-Hfl coli and incubated at 37°C for 30 minutes. Soft LB agarose was added at 48°C, mixed and poured onto prewarmed 22x22 cm rectangular LB agar plates and incubated overnight at 37°C.

35 Plaque lifts were performed by chilling the plates at 4°C to harden the top agarose and prevent it from peeling,

marking a nylon or nitrocellulose filter on the surface contacting the plaques, laying the filter on the surface without trapped air bubbles, and leaving it for about a minute. A number of asymmetric dots were inserted with
5 Indian ink with a syringe and needle so that the ink soaked into the agar. The sheets were then peeled gently away, and laid plaque side up on two sheets of Whattman 3MM soaked in denaturing solution, and left for about 2 minutes. The sheets were then peeled away and immersed in
10 a standard neutralizing solution for 5 minutes, immersed in 5X SSC, air dried, and baked at 80°C under vacuum, for 2 hours.

The filters were prehybridized in 40% formamide, 5X SSC, 0.1 % SDS, 1X Denhardt, and 100 ng/ml denatured
15 salmon sperm DNA at 37°-42°C for 1 hour. Labeled DNA probe (1 million cpm/ml) was denatured by heating at 100°C for 10 minutes, chilled, and then added to the prehybridization solution, and hybridized at 37°-42°C overnight. The filters were washed in 2X SSC and, 0.1%
20 SDS at 42°C or higher temperature.

Positive plaques were identified and purified by rescreening two more times. The probe was labeled by nick-translation.

Phage stocks were made by isolating and removing a
25 well separated plaque with the narrow end of an autoclaved Pasteur pipette, immersing it in 1 ml of standard SM buffer, and adding a drop of chloroform. This was left for a few hours at room temperature (20°C-24°C) or overnight at 4°C, vortexed, and centrifuged.

30 The cDNA insert was amplified by polymerase chain reactions (PCR). 20 microliters of phage stock was used in 100 microliters of standard PCR reaction buffer, by adding all components except Polymerase. This mixture was heated to 99°C, and Vent DNA polymerase (Biolabs) was
35 added to start the PCR cycles. The PCR conditions were 95°C 1 minute, 72°C 1 minute, 72°C 3 minutes (1 minute per

kilobase) for 30 cycles, 72°C 5 minutes, and kept at 4°C till further utilized.

The applicant isolated a clone from the cDNA library using an EcoR1-BglII fragment (nucleotides 450-909) of the
5 rat PPAR (459 bases) as a probe and the hybridization conditions provided above. This clone was purified and its sequence defined. This sequence is shown in Figure 1, and as SEQ. ID. NO. 1. Figure 2 is a comparison of mPPAR and hPPAR1 amino acid sequences with those amino acids
10 having identity between the two sequences enclosed in blocks.

Example 2: Northern blot analysis

A human multiple tissue Northern blot was purchased from Clontech. Screening was done following the
15 manufacturer's protocol. The blot was prehybridized in 5X SSPE, 10X Denhardt's solution, 100µg/ml of freshly denatured salmon sperm DNA, 50% formamide and 2% SDS for 3 hours at 42°C. DNA from the EcoR1 site at position 1025 of the coding region to the end of the cloned gene was
20 used as probe (see Figure 1). This DNA was labeled by random priming, boiled and added at a concentration of 1 million cpm/ml of prehybridization solution. Hybridization was carried out for 13 hours at 42°C. The blot was then washed in 2X SSC, 0.05% SDS at room
25 temperature followed by two washes in 0.1X SSC, 0.1% SDS at 50°C and exposed to X-ray film.

A specific band of about 10 kilobase was observed in all tissues except the brain. Maximal expression was observed in skeletal muscle, followed by heart, placenta,
30 pancreas, liver, kidney, and lung. The expression of hPPAR1 gene is therefore observed in tissues known to express PPARs in other species.

SEQUENCE LISTING

(1) GENERAL INFORMATION:

(i) APPLICANT:

5 (A) NAME: LIGAND PHARMACEUTICALS, INC.
(B) STREET: 9393 Towne Centre Drive
(C) CITY: San Diego
(D) STATE: California
(E) COUNTRY: United States of America
(F) POSTAL CODE (ZIP): 92121
10 (G) TELEPHONE: (619) 535-3900
(H) TELEFAX: (619) 535-3906

(ii) TITLE OF INVENTION: HUMAN PEROXISOME
PROLIFERATOR
15 ACTIVATED RECEPTOR

(iii) NUMBER OF SEQUENCES: 3

(iv) COMPUTER READABLE FORM:

20 (A) MEDIUM TYPE: 3.5" Diskette, 1.44 Mb
(B) COMPUTER: IBM compatible
(C) OPERATING SYSTEM: IBM P.C. DOS
(Version 5.0)
(D) SOFTWARE: WordPerfect (Version 5.1)

25 (v) CURRENT APPLICATION DATA:

APPLICATION NUMBER: To Be Assigned

(vi) PRIOR APPLICATION DATA:

30 (A) APPLICATION NUMBER: 08/141,500
(B) FILING DATE: 22-OCT-1993

(vi) PRIOR APPLICATION DATA:

(A) APPLICATION NUMBER: 08/143,215
(B) FILING DATE: 26-OCT-1993

35

(2) INFORMATION FOR SEQ ID NO: 1:

(i) SEQUENCE CHARACTERISTICS:

40 (A) LENGTH: 1407 base pairs
(B) TYPE: nucleic acid
(C) STRANDEDNESS: single
(D) TOPOLOGY: linear

(ii) SEQUENCE DESCRIPTION : SEQ ID NO: 1:

	ATG GTG GAC ACG GAA AGC CCA CTC TGC CCC CTC TCC CCA	39
	Met Val Asp Thr Glu Ser Pro Leu Cys Pro Leu Ser Pro	
	5 10	
5	CTC GAG GCC GGC GAT CTA GAG AGC CCG TTA TCT GAA GAG	78
	Leu Glu Ala Gly Asp Leu Glu Ser Pro Leu Ser Glu Glu	
	15 20 25	
	TTC CTG CAA GAA ATG GGA AAC ATC CAA GAG ATT TCG CAA	117
10	Phe Leu Gln Glu Met Gly Asn Ile Gln Glu Ile Ser Gln	
	30 35	
	TCC ATC GGC GAG GAT AGT TCT GGA AGC TTT GGC TTT ACG	156
	Ser Ile Gly Glu Asp Ser Ser Gly Ser Phe Gly Phe Thr	
	40 45 50	
15	GAA TAC CAG TAT TTA GGA AGC TGT CCT GGC TCA GAT GGC	195
	Glu Tyr Gln Tyr Leu Gly Ser Cys Pro Gly Ser Asp Gly	
	55 60 65	
	TCG GTC ATC ACG GAC ACG CTT TCA CCA GCT TCG AGC CCC	234
	Ser Val Ile Thr Asp Thr Leu Ser Pro Ala Ser Ser Pro	
	70 75	
20	TCC TCG GTG ACT TAT CCT GTG GTC CCC GGC AGC GTG GAC	273
	Ser Ser Val Thr Tyr Pro Val Val Pro Gly Ser Val Asp	
	80 85 90	
	GAG TCT CCC AGT GGA GCA TTG AAC ATC GAA TGT AGA ATC	312
25	Glu Ser Pro Ser Gly Ala Leu Asn Ile Glu Cys Arg Ile	
	95 100	
	TGC GGG GAC AAG GCC TCA GGC TAT CAT TAC GGA GTC CAC	351
	Cys Gly Asp Lys Ala Ser Gly Tyr His Tyr Gly Val His	
	105 110 115	
30	GCG TGT GAA GGC TGC AAG GGC TTC TTT CGG CGA ACG ATT	390
	Ala Cys Glu Gly Cys Lys Gly Phe Phe Arg Arg Thr Ile	
	120 125 130	
	CGA CTC AAG CTG GTG TAT GAC AAG TGC GAC CGC AGC TGC	429
	Arg Leu Lys Leu Val Tyr Asp Lys Cys Asp Arg Ser Cys	
	135 140	
35	AAG ATC CAG AAA AAG AAC AGT TTC AAA TGC CAG TAT TGT	468
	Lys Ile Gln Lys Lys Asn Arg Asn Lys Cys Gln Tyr Cys	
	145 150 155	
40	CGA TTT CAC AAG TGC CTT TCT GTC GGG ATG TCA CAC AAC	507
	Arg Phe His Lys Cys Leu Ser Val Gly Met Ser His Asn	
	160 165	

	GCG	ATT	CGT	TTT	GGA	CGA	ATG	CCA	AGA	TCT	GAG	AAA	GCA	546
	Ala	Ile	Arg	Phe	Gly	Arg	Met	Pro	Arg	Ser	Glu	Lys	Ala	
	170					175					180			
	AAA	CTG	AAA	GCA	GAA	ATT	CTT	ACC	TGT	GAA	CAT	GAC	ATA	585
5	Lys	Leu	Lys	Ala	Glu	Ile	Leu	Thr	Cys	Glu	His	Asp	Ile	
			185					190					195	
	GAA	GAT	TCT	GAA	ACT	GCA	GAT	CTC	AAA	TCT	CTG	GCC	AAG	624
	Glu	Asp	Ser	Glu	Thr	Ala	Asp	Leu	Lys	Ser	Leu	Ala	Lys	
					200					205				
10	AGA	ATC	TAC	GAG	GCC	TAC	TTG	AAG	AAC	TTC	AAC	ATG	AAC	663
	Arg	Ile	Tyr	Glu	Ala	Tyr	Leu	Lys	Asn	Phe	Asn	Met	Asn	
		210					215					220		
	AAG	GTC	AAA	GCC	CGG	GTC	ATC	CTC	TCA	GGA	AAG	GCC	AGT	702
15	Lys	Val	Lys	Ala	Arg	Val	Ile	Leu	Ser	Gly	Lys	Ala	Ser	
				225					230					
	AAC	AAT	CCA	CCT	TTT	GTC	ATA	CAT	GAT	ATG	GAG	ACA	CTG	741
	Asn	Asn	Pro	Pro	Phe	Val	Ile	His	Asp	Met	Glu	Thr	Leu	
	235					240					245			
	TGT	ATG	GCT	GAG	AAG	ACG	CTG	GTG	GCC	AAG	CTG	GTG	GCC	780
20	Cys	Met	Ala	Glu	Lys	Thr	Leu	Val	Ala	Lys	Leu	Val	Ala	
			250					255					260	
	AAT	GGC	ATC	CAG	AAC	AAG	GAG	GCG	GAG	GTC	CGC	ATC	TTT	819
	Asn	Gly	Ile	Gln	Asn	Lys	Glu	Ala	Glu	Val	Arg	Ile	Phe	
					265					270				
25	CAC	TCG	TGC	CAG	TGC	ACG	TCA	GTG	GTG	ACC	GTC	ACG	GAG	858
	His	Cys	Cys	Gln	Cys	Thr	Ser	Val	Glu	Thr	Val	Thr	Glu	
		275					280					285		
	CTC	ACG	GAA	TTC	GCC	AAG	GCC	ATC	CCA	GGC	TTC	GCA	AAC	897
30	Leu	Thr	Glu	Phe	Ala	Lys	Ala	Ile	Pro	Gly	Phe	Ala	Asn	
				290					295					
	TTG	GAC	CTG	AAC	GAT	CAA	GTG	ACA	TTG	CTA	AAA	TAC	GGA	936
	Leu	Asp	Leu	Asn	Asp	Gln	Val	Thr	Leu	Leu	Lys	Tyr	Gly	
	300					305					310			
	GTT	TAT	GAG	GCC	ATA	TTC	GCC	ATG	CTG	TCT	TCT	GTG	ATG	975
35	Val	Tyr	Glu	Ala	Ile	Phe	Ala	Met	Leu	Ser	Ser	Val	Met	
			315					320					325	
	AAC	AAA	GAC	GGG	ATG	CTG	GTA	GCG	TAT	GGA	AAT	GGG	TTT	1014
	Asn	Lys	Asp	Gly	Met	Leu	Val	Ala	Tyr	Gly	Asn	Gly	Phe	
					330					335				
40	ATA	ACT	CGT	GAA	TTC	CTA	AAA	AGC	CTA	AGG	AAA	CCG	TTC	1053
	Ile	Thr	Arg	Glu	Phe	Leu	Lys	Ser	Leu	Arg	Lys	Pro	Phe	
		340					345					350		

10

	TGT GAT ATC ATG GAA CCC AAG TTT GAT TTT GCC ATG AAG	1092
	Cys Asp Ile Met Glu Pro Lys Phe Asp Phe Ala Met Lys	
	355 360	
5	TTC AAT GCA CTG GAA CTG GAT GAC AGT GAT ATC TCC CTT	1131
	Phe Asn Ala Leu Glu Leu Asp Asp Ser Asp Ile Ser Leu	
	365 370 375	
	TTT GTG GCT GCT ATC ATT TGC TGT GGA GAT CGT CCT GGC	1170
	Phe Val Ala Ala Ile Ile Cys Cys Gly Asp Arg Pro Gly	
	380 385 390	
10	CTT CTA AAC GTA GGA CAC ATT GAA AAA ATG CAG GAG GGT	1209
	Leu Leu Asn Val Gly His Ile Glu Lys Met Gln Glu Gly	
	395 400	
	ATT GTA CAT GTG CTC AGA CTC CAC CTG CAG AGC AAC CAC	1248
	Ile Val His Val Leu Arg Leu His Leu Gln Ser Asn His	
15	405 410 415	
	CCG GAC GAT ATC TTT CTC TTC CCA AAA CTT CTT CAA AAA	1287
	Pro Asp Asp Ile Phe Leu Phe Pro Lys Leu Leu Gln Lys	
	420 425	
	ATG GCA GAC CTC CGG CAG CTG GTG ACG GAG CAT GCG CAG	1326
20	Met Ala Asp Leu Arg Gln Leu Val Thr Glu His Ala Gln	
	430 435 440	
	CTG GTG CAG ATC ATC AAG AAG ACG GAG TCG GAT CGT GCG	1365
	Leu Val Gln Ile Ile Lys Lys Thr Glu Ser Asp Ala Ala	
	445 450 455	
25	CTG CAC CCG CTA CTG CAG GAG ATC TAC AGG GAC ATG TAC	1404
	Leu His Pro Leu Leu Gln Glu Ile Tyr Arg Asp Met Tyr	
	460 465	
	TGA	1407

(2) INFORMATION FOR SEQ ID NO: 2:

30 (i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 468 amino acids
 (B) TYPE: amino acid
 (D) TOPOLOGY: linear

35 (ii) SEQUENCE DESCRIPTION : SEQ ID NO: 2

Met Val Asp Thr Glu Ser Pro Leu Cys Pro Leu Ser Pro
 5 10

Leu Glu Ala Gly Asp Leu Glu Ser Pro Leu Ser Glu Glu
 15 20 25

11

	Phe	Leu	Gln	Glu	Met	Gly	Asn	Ile	Gln	Glu	Ile	Ser	Gln
				30					35				
	Ser	Ile	Gly	Glu	Asp	Ser	Ser	Gly	Ser	Phe	Gly	Phe	Thr
	40					45					50		
5	Glu	Tyr	Gln	Tyr	Leu	Gly	Ser	Cys	Pro	Gly	Ser	Asp	Gly
			55					60					65
	Ser	Val	Ile	Thr	Asp	Thr	Leu	Ser	Pro	Ala	Ser	Ser	Pro
					70					75			
10	Ser	Ser	Val	Thr	Tyr	Pro	Val	Val	Pro	Gly	Ser	Val	Asp
		80					85					90	
	Glu	Ser	Pro	Ser	Gly	Ala	Leu	Asn	Ile	Glu	Cys	Arg	Ile
				95					100				
	Cys	Gly	Asp	Lys	Ala	Ser	Gly	Tyr	His	Tyr	Gly	Val	His
	105					110					115		
15	Ala	Cys	Glu	Gly	Cys	Lys	Gly	Phe	Phe	Arg	Arg	Thr	Ile
			120					125					130
	Arg	Leu	Lys	Leu	Val	Tyr	Asp	Lys	Cys	Asp	Arg	Ser	Cys
					135					140			
20	Lys	Ile	Gln	Lys	Lys	Asn	Arg	Asn	Lys	Cys	Gln	Tyr	Cys
	145					150						155	
	Arg	Phe	His	Lys	Cys	Leu	Ser	Val	Gly	Met	Ser	His	Asn
				160					165				
	Ala	Ile	Arg	Phe	Gly	Arg	Met	Pro	Arg	Ser	Glu	Lys	Ala
	170					175					180		
25	Lys	Leu	Lys	Ala	Glu	Ile	Leu	Thr	Cys	Glu	His	Asp	Ile
			185					190					195
	Glu	Asp	Ser	Glu	Thr	Ala	Asp	Leu	Lys	Ser	Leu	Ala	Lys
					200					205			
30	Arg	Ile	Tyr	Glu	Ala	Tyr	Leu	Lys	Asn	Phe	Asn	Met	Asn
	210						215					220	
	Lys	Val	Lys	Ala	Arg	Val	Ile	Leu	Ser	Gly	Lys	Ala	Ser
				225					230				
	Asn	Asn	Pro	Pro	Phe	Val	Ile	His	Asp	Met	Glu	Thr	Leu
	235					240					245		
35	Cys	Met	Ala	Glu	Lys	Thr	Leu	Val	Ala	Lys	Leu	Val	Ala
			250					255					260

12

	Asn	Gly	Ile	Gln	Asn	Lys	Glu	Ala	Glu	Val	Arg	Ile	Phe
					265					270			
	His	Cys	Cys	Gln	Cys	Thr	Ser	Val	Glu	Thr	Val	Thr	Glu
		275					280					285	
5	Leu	Thr	Glu	Phe	Ala	Lys	Ala	Ile	Pro	Gly	Phe	Ala	Asn
				290					295				
	Leu	Asp	Leu	Asn	Asp	Gln	Val	Thr	Leu	Leu	Lys	Tyr	Gly
	300					305					310		
10	Val	Tyr	Glu	Ala	Ile	Phe	Ala	Met	Leu	Ser	Ser	Val	Met
			315					320					325
	Asn	Lys	Asp	Gly	Met	Leu	Val	Ala	Tyr	Gly	Asn	Gly	Phe
					330					335			
	Ile	Thr	Arg	Glu	Phe	Leu	Lys	Ser	Leu	Arg	Lys	Pro	Phe
		340					345					350	
15	Cys	Asp	Ile	Met	Glu	Pro	Lys	Phe	Asp	Phe	Ala	Met	Lys
				355					360				
	Phe	Asn	Ala	Leu	Glu	Leu	Asp	Asp	Ser	Asp	Ile	Ser	Leu
	365					370					375		
20	Phe	Val	Ala	Ala	Ile	Ile	Cys	Cys	Gly	Asp	Arg	Pro	Gly
			380					385					390
	Leu	Leu	Asn	Val	Gly	His	Ile	Glu	Lys	Met	Gln	Glu	Gly
					395					400			
	Ile	Val	His	Val	Leu	Arg	Leu	His	Leu	Gln	Ser	Asn	His
		405					410					415	
25	Pro	Asp	Asp	Ile	Phe	Leu	Phe	Pro	Lys	Leu	Leu	Gln	Lys
				420					425				
	Met	Ala	Asp	Leu	Arg	Gln	Leu	Val	Thr	Glu	His	Ala	Gln
	430					435					440		
30	Leu	Val	Gln	Ile	Ile	Lys	Lys	Thr	Glu	Ser	Asp	Ala	Ala
			445					450					455
	Leu	His	Pro	Leu	Leu	Gln	Glu	Ile	Tyr	Arg	Asp	Met	Tyr
					460					465			468

13

(2) INFORMATION FOR SEQ ID NO: 3:

(i) SEQUENCE CHARACTERISTICS:

5 (A) LENGTH: 468 amino acids
 (B) TYPE: amino acid
 (D) TOPOLOGY: linear

(ii) SEQUENCE DESCRIPTION : SEQ ID NO: 3:

Met Val Asp Thr Glu Ser Pro Ile Cys Pro Leu Ser Pro
 5 10

10 Leu Glu Ala Asp Asp Leu Glu Ser Pro Leu Ser Glu Glu
 15 20 25

Phe Leu Gln Glu Met Gly Asn Ile Gln Glu Ile Ser Gln
 30 35

Ser Ile Gly Glu Glu Ser Ser Gly Ser Phe Gly Phe Ala
 40 45 50

15 Asp Tyr Gln Tyr Leu Gly Ser Cys Pro Gly Ser Glu Gly
 55 60 65

Ser Val Ile Thr Asp Thr Leu Ser Pro Arg Ser Ser Pro
 70 75

20 Ser Ser Val Ser Cys Pro Val Ile Pro Ala Ser Thr Asp
 80 85 90

Glu Ser Pro Gly Ser Ala Leu Asn Ile Glu Cys Arg Ile
 95 100

Cys Gly Asp Lys Ala Ser Gly Tyr His Tyr Gly Val His
 105 110 115

25 Ala Cys Glu Gly Cys Lys Gly Phe Phe Arg Arg Thr Ile
 120 125 130

Arg Leu Lys Leu Val Tyr Asp Lys Cys Asp Arg Ser Cys
 135 140

30 Lys Ile Gln Lys Lys Asn Arg Asn Lys Cys Gln Tyr Cys
 145 150 155

Arg Phe His Lys Cys Leu Ser Val Gly Met Ser His Asn
 160 165

35 Ala Ile Arg Phe Gly Arg Met Pro Arg Ser Glu Lys Ala
 170 175 180

Lys Leu Lys Ala Glu Ile Leu Thr Cys Glu His Asp Leu
 185 190 195

14

Lys Asp Ser Glu Thr Ala Asp Leu Lys Ser Leu Gly Lys
 200 205
 Arg Ile His Glu Ala Tyr Leu Lys Asn Phe Asn Met Asn
 210 215 220
 5 Lys Val Lys Ala Arg Val Ile Leu Ala Gly Lys Thr Ser
 225 230
 Asn Asn Pro Pro Phe Val Ile His Asp Met Glu Thr Leu
 235 240 245
 10 Cys Met Ala Glu Lys Thr Leu Val Ala Lys Met Val Ala
 250 255 260
 Asn Gly Val Glu Asp Lys Glu Ala Glu Val Arg Phe Phe
 265 270
 His Cys Cys Gln Cys Met Ser Val Glu Thr Val Thr Glu
 275 280 285
 15 Leu Thr Glu Phe Ala Lys Ala Ile Pro Gly Phe Ala Asn
 290 295
 Leu Asp Leu Asn Asp Gln Val Thr Leu Leu Lys Tyr Gly
 300 305 310
 20 Val Tyr Glu Ala Ile Phe Thr Met Leu Ser Ser Leu Met
 315 320 325
 Asn Lys Asp Gly Met Leu Ile Ala Tyr Gly Asn Gly Phe
 330 335
 Ile Thr Arg Glu Phe Leu Lys Asn Leu Arg Lys Pro Phe
 340 345 350
 25 Cys Asp Ile Met Glu Pro Lys Phe Asp Phe Ala Met Lys
 355 360
 Phe Asn Ala Leu Glu Leu Asp Asp Ser Asp Ile Ser Leu
 365 370 375
 30 Phe Val Ala Ala Ile Ile Cys Cys Gly Asp Arg Pro Gly
 380 385 390
 Leu Leu Asn Ile Gly Tyr Ile Glu Lys Leu Gln Glu Gly
 395 400
 Ile Val His Val Leu Lys Leu His Leu Gln Ser Asn His
 405 410 415
 35 Pro Asp Asp Thr Phe Leu Phe Pro Lys Leu Leu Gln Lys
 420 425

15

Met Val Asp Leu Arg Gln Leu Val Thr Glu His Ala Gln
430 435 440

Leu Val Gln Val Ile Lys Lys Thr Glu Ser Asp Ala Ala
445 450 455

5 Leu His Pro Leu Leu Gln Glu Ile Tyr Arg Asp Met Tyr
460 465 468

What is claimed is:

1. Purified nucleic acid comprising the nucleotide sequence shown in SEQ ID NO. 1.
2. A vector comprising said nucleic acid of claim
5 1.
3. Recombinant PPAR expressed from said nucleic acid of claim 1.

1/2

10 20 30 40 50 60 70 80 90 100
 1234567890 1234567890 1234567890 1234567890 1234567890 1234567890 1234567890 1234567890 1234567890 1234567890
 ATGGTGGACA CGGAAGGCC ACTCTGCC CTTCTCCAC TCTGCGCGG CGATCTAGG AGCCGTTAT CTGAAGATT CTGCAAGAA ATGGGAACA
 M V D T E S P L C P L S P L E A G D L E S P L S E E F L Q E M G N I
 200
 TCAAGAGAT TTCGAATCC ATCGGCGAGG ATAGTTCTGG AAGCTTTGGC TTACGGAAT ACCAGTATT AGGAAGCTGT CTGGCTCAG ATGGCTCGGT
 Q E I S Q S I G E D S S G S F G F T E Y Q Y L G S C P G S D G S V
 300
 CATCAGGAC ACGTTTTCAC CAGCTTCGAG CCCCTCCTCG GTGACTTATC CTGTGGTCCC CGGAGCGTG GACGAGTCTC CCAGTGGAGC ATTGAACATC
 I T D T L S P A S S P S S V T Y P V V P G S V D E S P S G A L N I
 400
 GAATGTAGAA TCTGCGGGA CAAGGCTCA GGTATCAT TACGAGTCCA CCGGTGTGAA GCTTCITTCG GCGAAGCATT CGACTCAAGC
 E C R I C G D K A S G Y H Y G V H A C E G C K G F F R R T I R L K L
 500
 TGGTGTATGA CAATGCGAC CGCAGCTGCA AGATCCAGAA AAAGAACAGA AACAAATGCC AGTATTGTGC ATTTACAAG TGCCTTCTG TCGGGATGC
 V Y D K C D R S C K I Q K K N R N K C Q Y C R F H K C L S V G M S
 600
 ACACAGCGG ATTCTTTTG GACGAATGCC AAGATCTGAG AAGCAAAAC TGAAGCAGA AATTCTTACC TGTGAACATG ACATAGAAGA TTCTGAAACT
 H N A I R F G R M P R S E K A K L K A E I L T C E H D I E D S E T
 700
 GCAGATCTCA AATCTCTGGC CAAGAGAATC TACGAGGCT ACTTGAAGAA CTTCACATG AACCAAGTCA AAGCCGGGT CATCTCTCA GGAAGGCCA
 A D L K S L A K R I Y E A Y L K N F N M N K V K A R V I L S G K A S
 800
 GTAACAATCC ACCTTTTGTG ATACATGATA TGGAGACACT GTGTATGGCT GAGAAGACG TGGTGGCCAA GCTGTGGCC AATGGCATCC AGAACAAGGA
 N N P P F V I H D M E T L C M A E K T L V A K L V A N G I Q N K E
 900
 GCGGAGGTC CGCATCTTTC ACTGCTGCA GTGCAGTCA GTGGAGACCG TCAGGAGCT CACGGAATTC GCCAAGGCCA TCCCAGGCTT CGCAAACTTG
 A E V R I F H C C Q C T S V E T V T E L T E F A K A I P G F A N L
 1000
 GACCTGAACG ATCAAGTGAC ATTGCTAAAA TACGAGTTT ATGAGGCCAT ATTGCGCATG CTGTCTCTG TGATGAACAA AGACGGGATG CTGTAGCGT
 D L N D Q V T L L K Y G V Y E A I F A M L S S V M N K D G M L V A Y
 1100
 ATGGAATGG GTTTATAACT CGTGAATTC TAAAAAGCCT AAGGAAACCG TTCTGTGATA TCATGGAACC CAAGTTTGT TTTGCCATGA AGTTCAATGC
 G N G F I T R E F L K S L R K P F C D I M E P K F D F A M K F N A
 1200
 ACTGGAACG GATGACAGTG ATATCTCCCT TTTTGTGGCT GCTATCATTT GCTGTGAGA TCGTCTCTGC CTCTAAACG TAGGACACAT TGA AAAAATG
 L E L D D S D I S L F V A A I I C C G D R P G L L N V G H I E K M
 1300
 CAGGAGGTA TTGTACATGT GCTCAGACT CACCTGCAGA GCAACCAACC GGACGATATC TTCTCTTCC CAAAATTCT TCAAAAATG GCAGACTCC
 Q E G I V H V L R L H L Q S N H P D D I F L F P K L L Q K M A D L R
 1400
 GGCAGCTGGT GACGAGCAT GCGCAGCTGG TGCAGATCAT CAAGAAGACG GAGTGGGATG CTGCGCTGCA CCGCTACTG CAGGAGATCT ACAGGACAT
 Q L V T E H A Q L V Q I I K K T E S D A A L H P L L Q E I Y R D M
 1407
 GTACTGA
 Y X

FIG. 1

RECTIFIED SHEET (RULE 91)

ISA/EP

2/2

MVDTESRCP	LSPLEADLE	SPLSEEFLOE	MGNIQEISQS	IGESGSGFG	FADYQYLGSC	PGSEGSVITD	TLSPRSSPSS	VSCPVIPAST	DESPGSALNI	100
MVDTESRCP	LSPLEADLE	SPLSEEFLOE	MGNIQEISQS	IGESGSGFG	FADYQYLGSC	PGSEGSVITD	TLSPRSSPSS	VSCPVIPAST	DESPGSALNI	100
ECRICGDKAS	GYHYGVHACE	GCKGFFRRTI	RKCLVYDKCD	RSCKIQKKNR	NKCQYCRFHK	CLSVGMSHNA	IRFGMRPRSE	KAKLKAELT	CEHDLKDESET	200
ECRICGDKAS	GYHYGVHACE	GCKGFFRRTI	RKCLVYDKCD	RSCKIQKKNR	NKCQYCRFHK	CLSVGMSHNA	IRFGMRPRSE	KAKLKAELT	CEHDLKDESET	200
ADLKSJGKRI	YEAYLKNFNM	NKVAKRVILA	GKISNNPPFV	IHDMETLCMA	EKTLVAKMVA	NGVEDKEAEV	RHFHCCQQMS	VETVTELTEF	AKAIPGFANL	300
ADLKSJGKRI	YEAYLKNFNM	NKVAKRVILS	GKISNNPPFV	IHDMETLCMA	EKTLVAKMVA	NGIQNKEAEV	RHFHCCQQIS	VETVTELTEF	AKAIPGFANL	300
DLNDQVTLK	YGVEAIFIM	LSSMKNKOGM	LIAYGNGFIT	REFLNLRLKP	FCDIMEPKFD	FAMKFNALEL	DDSDISLFVA	AIICCGDRPG	LLNIGMIELK	400
DLNDQVTLK	YGVEAIFAM	LSSMKNKOGM	LIAYGNGFIT	REFLNLRLKP	FCDIMEPKFD	FAMKFNALEL	DDSDISLFVA	AIICCGDRPG	LLNIGMIELM	400
QEGIVHVLL	HLQSNHPDDI	FLFPKLLQKM	VDLRQLVTEH	AQLVQMIIKKT	ESDAALHPLL	QEIYRDMY-				468
QEGIVHVLL	HLQSNHPDDI	FLFPKLLQKM	ADLRQLVTEH	AQLVQMIIKKT	ESDAALHPLL	QEIYRDMYX				469

FIG. 2